

# COVID-19 Vaccines Currently Available at the MTPL (Updated 10/22/21)

Pfizer	Moderna	J&J
1 <sup>st</sup> dose	1 <sup>st</sup> dose	1 dose
2 <sup>nd</sup> dose	2 <sup>nd</sup> dose	Booster – everyone ages 18 and older who received a first Johnson & Johnson dose may receive a booster dose at least two months after completing the single-dose primary regimen.
3 <sup>rd</sup> full dose/booster (same ml) at least 6 months from 2 <sup>nd</sup> dose for: <ul style="list-style-type: none"> <li>• individuals 65 and older</li> <li>• Long-Term Care residents</li> <li>• Anyone 18-64 years of age with underlying medical conditions</li> <li>• Individuals who work in healthcare settings or an occupation that makes them high-risk for contracting COVID</li> </ul>	3 <sup>rd</sup> full dose for those who have: <ul style="list-style-type: none"> <li>• Been receiving active cancer treatment for tumors or cancers of the blood</li> <li>• Received an organ transplant and are taking medicine to suppress the immune system</li> <li>• Received a stem cell transplant within the last 2 years or are taking medicine to suppress the immune system</li> <li>• Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome)</li> <li>• Advanced or untreated HIV infection</li> <li>• Active treatment with high-dose corticosteroids or other drugs that may suppress your immune response</li> </ul>	
	Booster (half dose) – six months or more after their initial series for: <ul style="list-style-type: none"> <li>• 65 years and older</li> <li>• Age 18+ who live in long-term care settings</li> <li>• Age 18+ who have underlying medical conditions</li> <li>• Age 18+ who work or live in high-risk settings</li> </ul>	